



United Phosphorus, Inc.  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406

TEBUZOL™ 3.6 F Fungicide  
For Use on Wheat to Suppress Fusarium head  
blight (scab)  
SECTION 18 SPECIFIC EXEMPTION  
File Symbol: 07MT04

## FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MONTANA UNDER SECTION 18 SPECIFIC EXEMPTION

These use directions must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its use directions. Read entire Directions For Use and Conditions of Sale and Limitation of Warranty and Liability before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for TEBUZOL 3.6 F Fungicide (EPA Reg. No. 70506-114). Any adverse effects resulting from the use of TEBUZOL 3.6 F Fungicide under this specific exemption must be immediately reported to the Montana Department of Agriculture.

**EFFECTIVE PERIOD:** Expires on July 20, 2007

### DIRECTIONS FOR USE

#### RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF TEBUZOL 3.6 F
Wheat (including durum)	<i>Fusarium</i> head blight (scab)	4 fl. oz. per acre

Apply TEBUZOL 3.6 F as a preventive foliar spray. For optimum disease suppression, the lowest rate of a spray surfactant should be tank-mixed with TEBUZOL 3.6 F. TEBUZOL 3.6 F may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 3 gallons of spray solution per acre by aircraft spray equipment.

Application through any type of irrigation equipment is prohibited.

Do not make more than 1 application of a tebuconazole-containing product to wheat per season. One application may be made by ground or air at a rate of 4 fl. oz. product per acre per year. A maximum of 300,500 acres of wheat in Montana may be treated. TEBUZOL 3.6 F may be applied up through the beginning of flowering (Feekes 10.51). Application may not be made within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Do not enter treated field for 12 hours after application.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**For Emergency Medical Assistance,  
call the National Pesticide Information Center 1-800-858-7378.  
For chemical emergency: spill, leak, fire, exposure, or accident,  
call CHEMTREC 1-800-424-9300.**